

ACT Treatment Protocol

ENVIRONMENTAL EMERGENCIES – COLD EXPOSURE

When cold exposure affects the entire body: hypothermia or general cooling develops. When cold exposure affects a particular body part: local cooling, or frostbite occurs. Frostbite most commonly affects the ears, nose, face, hands, feet, and toes.

- A. Perform **Initial Treatment / Universal Patient Care Protocol** and follow the proper protocol for medical management based on clinical presentation.
 - 1. Place patient in warm environment.
 - 2. Treat with warm, humidified oxygen and warmed IV fluids.
 - 3. Remove all wet clothing.
 - 4. Insulate core (head, neck, and trunk) with warm blankets.
 - 5. Rapid smooth transport.
- B. If patient is hypothermic, alert, and responding appropriately:
 - 1. Keep the patient still and handle very gently.
 - 2. Actively rewarm the patient by applying heat packs, hot water bottles, or electric heating pads to neck, chest, and abdomen.
 - 3. Allow patient to slowly drink warm fluids, but do not allow patient to drink stimulants.
 - 4. In **consultation with Medical Command**, establish mode (ground vs. air) and destination of transport.



- 5. Monitor vital signs closely during transport.
- C. If patient is hypothermic, unconscious or not responding appropriately:
 - 1. Handle patient as gently as possible and expedite transport.
 - 2. Wrap patient in insulated blankets for passive rewarming only.
 - 3. Give nothing by mouth.
 - 4. Continue IV normal saline at KVO.
 - 5. If patient has no pulse, perform CPR with the following cautions:



ACT Treatment Protocol

ENVIRONMENTAL EMERGENCIES – COLD EXPOSURE

- a. Check pulse for at least 60 seconds.
- b. Defibrillate VF/VT at max joules.
- c. Withhold IV medications until patient is rewarmed to core temperature of > 86° F.
- 6. Expedite transport.
- 7. In **consultation with Medical Command**, establish mode (ground vs. air) and destination of transport.



8. Further treatment per **order of Medical Command.**

D. Frostbite:

- 1. Remove constrictive clothing and jewelry and cover with dry dressing.
- 2. **Do not** rub, massage area or break blisters. Do not apply direct heat, allow patient to use affected area, or re-expose to cold.
- 3. Transport and notify **Medical Command**.